

## **Managing Strong Emotional Reactions to Traumatic Events: Tips for Families and Teachers**

When a large-scale tragedy occurs, it can cause strong and deeply felt reactions in adults and children.

***How adults express their emotions will influence the reactions of children and youth.***

Caregivers can help youngsters manage their feelings by modeling healthy coping strategies and closely monitoring their own emotional state and that of those in their care.

### **Common Reactions to Trauma**

It is not uncommon to feel any or all of the following:

- Shock/disbelief
- Fear
- Guilt
- Grief
- Confusion
- Shame/loss
- Anger

These reactions are often closely linked and can be difficult to separate, (e.g., where does grief end and outrage begin). Children, in particular, may have trouble understanding and talking about their feelings. Emotional reactions take place over a period of time and may not happen in any particular order. They can affect our behavior, our ability to function, and our overall sense of well-being. The intensity and ways we express our reactions will vary depending on our personal experience, general mental health, other stress factors in our lives, our coping style, our ability to self-monitor our emotional state, and our support network. Children of all ages may need guidance and support from the significant adults in their lives as they work through their thoughts and emotional reactions to the event.

### **Anger: A Natural Reaction**

For many people, adults and children alike, anger will be a natural extension of other emotions because it is a defensive mechanism that makes us *feel* more in control. As well, anger with the perpetrators of these horrible acts is, in many ways, justified. The desire to “retaliate” can be strong—but quite harmful if not redirected to a positive outlet. In some situations, a significant factor in the expression of anger is when there is an “enemy” on which to focus our feelings with whom others might be inappropriately associated. Adults must ensure that children do not “take out” their anger in inappropriate ways, such as lashing out at classmates or neighbors who might be unfairly associated with the perpetrators of violence because of their ethnicity or other affiliations. The key is to direct anger and other strong emotions in socially and psychologically healthy ways.

### **Recognizing Anger**

The first step in helping children manage their anger is getting your anger under control. Be aware of cues in your own behavior. If necessary, ask someone you trust (a family member, friend or colleague) to give you feedback on your anger reaction, and seek help, if needed. Observe the behavior of other adults around you and your child(ren) and be supportive if they show signs of increased anger.

## **Signs of Anger in Adults**

- Short temper/impatience
- Sleep and/or eating problems
- Restlessness and agitation
- Hitting and slamming objects, pets, or people
- Desire to inflict harm
- Verbal outbursts toward family, friends, or fellow workers
- A sense of losing control over your life
- Poor concentration or attention span
- Obsessing about the event
- Physical health affected; increase in blood pressure, dizzy, headaches, heart rate elevated, clenched jaw, knot in the stomach, and tight muscles, etc.
- You feel life should be fair, but it is not; and things are not how you want them to be
- Social media posts expressing intolerance and/or anger

## **Signs of Anger in Children**

- Behavioral outbursts, many times without an obvious cause
- Sleep problems
- Fights at school or home
- Physical attacks on others or animals
- Disobedience from otherwise well behaved child(ren)
- Child states he/she is really sad and does not know why
- Complaints of stomachaches and headaches; or vague aches and pains
- Other reactions similar to those of adults
- Social media posts expressing intolerance and/or anger

## **Dealing With Anger**

Some people have more problems than others dealing with anger. They either try to deny or ignore their feelings and keep them inside, or overreact and “blow-up.” These negative coping strategies can be physically and emotionally unhealthy. Pretending we don’t feel badly can have long-term effects that may eventually cause us to “lose it.” Conversely, psychological research shows that acting out your anger will not relieve it, but instead will make it more intense. We can learn to control or diffuse anger by how we think about the event or people involved and by finding other ways to regain our sense of control and security. Anger can also mask other emotions, such as grief, loss, or fear. It is important to address these related emotions as a way to deal with angry feelings.

## **Controlling Your Anger**

- Admit you are angry.
- Recognize this is a common reaction to an overwhelming event. It is how you control and manage your reaction that makes the difference.
- Try to identify the related emotions that may be fueling your anger, (e.g., sorrow, fear.)
- Find appropriate outlets for these related and equally important emotions, (e.g., talk with family members or friends, seek grief counseling, get involved in activities to help victims, etc.)
- Understand that *how* you think about the event can unnecessarily contribute to anger and other strong emotions. Develop a “positive” outlook and look for what can be done to help rather than harm.
- Stop, take a deep breath, visualize something peaceful or enjoyable, and try to relax for a few minutes.
- Avoid/decrease negative ways to cope, such as misuse of alcohol or drugs.
- Find an acceptable outlet, such as exercise, getting involved with your favorite hobby, sports.
- Distract yourself from continuing to think about the event -- call a friend, participate in an enjoyable activity.
- Keep a sense of humor.
- Turn off the TV and radio; play your favorite music.
- Maintain daily routines.

- Consult your doctor or a mental health specialist if your reactions continue to intensify, or you feel like doing harm to yourself or others.
- If you are seeing a mental health professional, be sure to share your angry feelings.

### **Helping Children Control Anger**

Realize they will imitate your responses and reactions.

- Let them understand anger is a normal emotion under these circumstances. However, acting out anger, hurting others, and uncontrolled anger is not okay.
- Answer their questions honestly and openly; but always consider their developmental age.
- Make family time to talk to the child(ren) about their reaction to the events.
- Highlight the many positive examples of people helping others and coming together as a community.
- Have child(ren) come up with ideas on how to help those who have been injured or otherwise effected by the tragedy.
- Teach them to stop, take a deep breath, and imagine a restful scene or enjoyable activity for a few minutes as a way to relax.
- Turn off the TV and make sure exposure to violence in the media, including social media, is restricted or monitored.
- Try to understand and encourage children to talk about their fears/sense of loss.
- Try to help them see how they would feel if someone hurt, yelled at, or hit them.
- Reinforce that the act of one person should not be generalized to an entire group. It is unfair to blame others who have no direct connection to the event.
- Sports, exercise, or other physical activity can be quite helpful.
- Be flexible in discipline and monitor your reactions to their misbehaviors.
- Seek mental health or physician consultation if these reactions do not clear up after 30 days.
- Keep family and school routines; get back to a normal life schedule as soon as possible.
- If age permits, get the child involved in volunteer work or community service, such as the Red Cross or Animal Shelter, where a child can feel that he/she is making a difference.

### **Warning Signs of Serious Emotional Trauma**

While strong emotional reactions to tragic events are normal, most will fade over the following weeks and most children soon will be able to resume normal activities with minimal displays of anger or anxiety. However, if any of the following symptoms or behaviors continue beyond a few weeks, or if any of these symptoms are exhibited to such a degree that it severely impacts the child's ability to participate in school or home activities, parents and teachers should seek mental health services for evaluation and possible treatment.

- Disruption in peer relationships (little or no interactions with friends, significant increase in conflict with classmates or friend).
- Strained family relationships (high degree of misbehavior, lashing out against family members, refusal to participate in normal family routines).
- Significant decrease in school performance.
- Ongoing physical complaints with no apparent cause.
- Use of alcohol or other drugs (or increase in comparison to previous behavior).
- Repeated nightmares and reporting strong fears of death, violence, etc.
- Repetitive play re-enacting the traumatic events.
- Low self esteem, negative talk about self (if this was not apparent prior to the trauma).
- General lack of energy and lack of interest in previously enjoyed activities.

Parents and teachers can help children overcome traumatic effects of a tragedy and use the process as an opportunity to teach them how to cope more effectively and deal with new challenges. Depending on the scope of the event, the process may take time and patience and the willingness to reach out to friends, neighbors, and co-workers to lend mutual support.